

Vanishing index for supersymmetric 2-matrix model with odd dimensional gauge group

G.M. Graf^(a), D. Hasler^(a), J. Hoppe^(b)

^(a) *Theoretische Physik, ETH-Hönggerberg, CH-8093 Zürich*

^(b) *Royal Institute of Technology, Department of Mathematics, S-10044 Stockholm*

February 1, 2008

Abstract

We define an operator which for odd-dimensional compact gauge group furnishes unitary equivalence of the bosonic and fermionic sector in the supersymmetric quantum-mechanical matrix model obtained by dimensional reduction from 3-dimensional supersymmetric Yang-Mills theory.

Let G be a compact semisimple Lie group. We consider a class of supersymmetric G -invariant quantum mechanical models involving $d = 2$ copies of the Lie algebra of G . If the dimension $g = \dim G$ of G is odd, we will show that the Hodge $*$ operator, in a simple, direct way, gives rise to a unitary equivalence of the Hamiltonian restricted to the bosonic and fermionic sector. This trivially implies the vanishing of the Witten-index (see [1-7] for work on the subject).

Let us first describe the Hilbert space. The configuration space of the bosonic degrees of freedom is $X = \mathbb{R}^{2g}$, with coordinates

$$(q_{sA})_{s=1,2, A=1,\dots,g}.$$

To describe the fermionic degrees of freedom one considers the Clifford algebra with hermitian generators $(\Theta_{\alpha A})_{\alpha=1,2, A=1,\dots,g}$ and commutation relations

$$\{\Theta_{\alpha A}, \Theta_{\beta B}\} = \delta_{\alpha\beta} \delta_{AB}.$$

We realize the Clifford algebra on the fermionic Fock space Λ over the Hilbert space $\mathbb{C}^g$. We decompose

$$\Lambda = \bigoplus_{p=0} \Lambda_p, \quad \Lambda_+ = \bigoplus_{p=0} \Lambda_{2p}, \quad \Lambda_- = \bigoplus_{p=0} \Lambda_{2p+1},$$

where $\Lambda_p = \bigwedge^p(\mathbb{C}^g)$ denotes the p -particle space. Λ carries a natural scalar product $\langle \cdot, \cdot \rangle$. Let

$$(e_A)_{A=1,\dots,g}$$

be the standard basis in $\mathbb{C}^g$, and $\lambda_A, \frac{\partial}{\partial \lambda_A}$ the associated creation and annihilation operators. The Clifford generators can now be expressed as

$$\Theta_{1A} = \frac{1}{\sqrt{2}} \left(\lambda_A + \frac{\partial}{\partial \lambda_A} \right), \quad \Theta_{2A} = \frac{i}{\sqrt{2}} \left(\lambda_A - \frac{\partial}{\partial \lambda_A} \right).$$

Let $(-1)^F = \left(\frac{i}{2}\right)^g \prod_A \Theta_{1A} \Theta_{2A}$, which equals ± 1 on $\Lambda_{\pm}$. We define the Hilbert spaces

$$\mathcal{H} = L^2(X) \otimes \Lambda = \mathcal{H}_+ \oplus \mathcal{H}_-,$$

with $\mathcal{H}_{\pm} = L^2(X) \otimes \Lambda_{\pm}$, called the bosonic and fermionic sector, respectively. The Hilbert space $\mathcal{H}$ carries a unitary representation of G . Let $\mathcal{H}^{(0)}$ denote the physical Hilbert space consisting of G -invariant states in $\mathcal{H}$. $\mathcal{H}_+^{(0)}, \mathcal{H}_-^{(0)}$ are given analogous. We define the Hodge $*$ operator with respect to the volume form $\omega = e_1 \wedge \dots \wedge e_g$ in Λ , i.e.

$$*: \Lambda \rightarrow \Lambda :$$

for $\beta \in \Lambda_p$, let $*\beta \in \Lambda_{m-p}$ be the unique element such that

$$\langle *\beta, \alpha \rangle = \langle \omega, \bar{\beta} \wedge \alpha \rangle, \text{ for all } \alpha \in \Lambda_{m-p}.$$

By $\bar{\beta}$ we mean complex conjugation of β . We list some properties of the Hodge $*$ in the following lemma, which can easily be verified.

Lemma 1

- (a) $*^{\dagger} = *^{-1}$
- (b) On Λ_p we have, $*^2 = (-1)^{(m-p)p}$.
- (c) If m is odd, then $*: \Lambda_+ \rightarrow \Lambda_-$ and $*: \Lambda_- \rightarrow \Lambda_+$, i.e. $\{*, (-1)^F\} = 0$.
- (d) On Λ_p we have, $\lambda_A * = (-1)^{p-1} * \frac{\partial}{\partial \lambda_A}$ and $\frac{\partial}{\partial \lambda_A} * = (-1)^p * \lambda_A$.

The model is defined by the following operators. Let f_{ABC} be the real and totally antisymmetric structure constants, with respect to an orthonormal basis of the Lie algebra of G . The generators of the unitary representation on $\mathcal{H}$ are

$$L_A = -if_{ABC} \left(q_{sB} \frac{\partial}{\partial q_{sC}} + \lambda_B \frac{\partial}{\partial \lambda_C} \right), \quad A = 1, \dots, g$$

The supercharges are given by

$$\begin{aligned} Q &= \left(\frac{\partial}{\partial q_{1A}} - i \frac{\partial}{\partial q_{2A}} \right) \frac{\partial}{\partial \lambda_A} + if_{ABC} q_{1A} q_{2B} \lambda_C, \\ Q^{\dagger} &= - \left(\frac{\partial}{\partial q_{1A}} + i \frac{\partial}{\partial q_{2A}} \right) \lambda_A - if_{ABC} q_{1A} q_{2B} \frac{\partial}{\partial \lambda_C} \end{aligned}$$

and the Hamiltonian by

$$H = -\Delta + \frac{1}{2} (f_{ABC} q_{mB} q_{nC})^2 + (q_{1A} + i q_{2A}) f_{ABC} \lambda_B \lambda_C - (q_{1A} - i q_{2A}) f_{ABC} \frac{\partial}{\partial \lambda_B} \frac{\partial}{\partial \lambda_C}.$$

The physical Hilbert space can be written as

$$\mathcal{H}^{(0)} = \{\psi \in \mathcal{H} \mid L_A \psi = 0, A = 1, \dots, g\}.$$

We recall the super-algebra

$$\begin{aligned} \{Q, Q^\dagger\}|_{\mathcal{H}^{(0)}} &= H|_{\mathcal{H}^{(0)}}, \quad \{Q, Q\}|_{\mathcal{H}^{(0)}} = 0, \quad \{Q^\dagger, Q^\dagger\}|_{\mathcal{H}^{(0)}} = 0, \\ \{(-1)^F, Q\} &= 0, \quad \{(-1)^F, Q^\dagger\} = 0. \end{aligned}$$

We define the following operators S and P by

$$\begin{aligned} S &: L^2(X) \rightarrow L^2(X) \\ \psi &\mapsto S\psi(q_{1A}, q_{2A}) = \psi(q_{1A}, -q_{2A}), \\ P &= S^* = *S. \end{aligned}$$

P has the following properties.

Theorem 2

- (a) P is unitary.
- (b) If $\dim G$ is odd, then we have $P^2 = 1$ and $P : \mathcal{H}_+ \rightarrow \mathcal{H}_-, P : \mathcal{H}_- \rightarrow \mathcal{H}_+$, i.e. $\{P, (-1)^F\} = 0$.
- (c) $[P, H] = 0$, $[P, L_A] = 0$.
- (d) On $\mathcal{H}_-$ we have $QP = PQ^\dagger$, $Q^\dagger P = -PQ$ and on $\mathcal{H}_+$, $QP = -PQ^\dagger$, $Q^\dagger P = PQ$.

Proof of Theorem.

- (a) follows by $S^\dagger = S^{-1}$ and Lemma 1(a).
- (b) follows from $S^{-1} = S$ and Lemma 1(b),(c).
- (c) Applying Lemma 1(d), we find

$$*\lambda_B \lambda_C = -\frac{\partial}{\partial \lambda_B} \frac{\partial}{\partial \lambda_C} *, \quad *\frac{\partial}{\partial \lambda_B} \frac{\partial}{\partial \lambda_C} = -\lambda_B \lambda_C *, \quad *\frac{\partial}{\partial \lambda_B} \lambda_C = \lambda_B \frac{\partial}{\partial \lambda_C} *,$$

which shows that $[P, H] = 0$ and $[P, L_A] = 0$.

- (d). Applying Lemma 1(d), we find

$$\begin{aligned} \text{on } \Lambda_+ : \quad \lambda_A * &= -*\frac{\partial}{\partial \lambda_A}, \quad \frac{\partial}{\partial \lambda_A} * = *\lambda_A, \\ \text{on } \Lambda_- : \quad \lambda_A * &= *\frac{\partial}{\partial \lambda_A}, \quad \frac{\partial}{\partial \lambda_A} * = -*\lambda_A, \end{aligned}$$

from which (c) follows. $\square$

This theorem directly implies the following corollary.

Corollary 3 *Let $\dim G$ be odd.*

- (a) $P : \mathcal{H}_+^{(0)} \rightarrow \mathcal{H}_-^{(0)}, P : \mathcal{H}_-^{(0)} \rightarrow \mathcal{H}_+^{(0)}$.
- (b) $P(H|_{\mathcal{H}_+^{(0)}})P = H|_{\mathcal{H}_-^{(0)}}$, i.e. $H|_{\mathcal{H}_+^{(0)}}$ and $H|_{\mathcal{H}_-^{(0)}}$ are unitarily aequivalent.

The corollary implies that the Witten index vanishes. In fact, both the principle and deficit contribution to the Witten index (cp. [2-7]) vanish, since P commutes with the characteristic function of the ball $B_R(0) = \{x : \|x\| \leq R\}$ for any R .

Acknowledgments. We thank J. Fröhlich for valuable discussions.

References

- [1] J. Fröhlich, J. Hoppe, On zero-mass ground states in super-membrane matrix models. Comm. Math. Phys. **191**, (1998) 613-626
- [2] P. Yi, Witten Index and threshold bound states of D-Branes, Nucl Phys. B **505** , (1997) 307
- [3] S. Sethi, M. Stern, Comm. Math. Phys. **194**, (1998) 675
- [4] W. Krauth, H. Nicolai, M. Staudacher, Phys. Lett. **B431**, (1998) 31
- [5] M.B. Green, M. Gutperle, D-particle bound states and the D-instanton measure, JHEP 9801 (1998) 005
- [6] G. Moore, N. Nekrasov, S. Shatashvili, D-particle bound states and generalized instantons, Comm. Math. Phys. **209**, (2000) 77-95
- [7] V.G. Kac, A.V. Smilga, Normalized Vacuum States in $N = 4$ Supersymmetric Yang–Mills Quantum Mechanics with Any Gauge Group, Nucl.Phys. B **571**, (2000) 515-554